ELSEVIER

Carbohydrate Research 300 (1997) XI–XIV

CARBOHYDRATE RESEARCH

Contents

Special Issue

Conformational Studies of Carbohydrates

Edited by E.F. Hounsell and M. Ragazzi (Guest Editor)

M. Ragazzi (Milan, Italy)	v	Guest Editor Introduction
	vii	Graphical Abstracts
J. Carver (Toronto, Canada)	1	Three-dimensional structures of carbohydrates and how they interact with proteins (from a transcript of the opening comments)
A. Poveda, J.L. Asensio, M. Martín-Pastor and J. Jiménez-Barbero (Madrid, Spain)	3	Solution conformation and dynamics of a tetrasaccharide related to the Lewis^X antigen deduced by ^{1}H NMR NOESY, ROESY, and T-ROESY measurements
E. Monteagudo, A. Madami, F. Animati, P. Lombardi and F. Arcamone (Pomezia, Rome, Italy)	11	Conformational analysis of 4-demethoxy-7-*O*-[2,6-dideoxy-4-*O*-(2,3,6-trideoxy-3-amino-α-L-*lyxo*-hexopyranosyl)-α-L-*lyxo*-hexopyranosyl]adriamicinone, the first doxorubicin disaccharide analogue to be reported
K. Ogawa (Osaka, Japan)	17	Progress in structure analyses on carbohydrates and polysaccharides
L. Catoire, C. Derouet, A.-M. Redon, R. Goldberg and C. Hervé du Penhoat (Paris, France)	19	An NMR study of the dynamic single-stranded conformation of sodium pectate
N.C. Santos, A.M.A. Sousa (Lisboa, Portugal), D. Betbeder (Labège, France), M. Prieto and M.A.R.B. Castanho (Lisboa, Portugal)	31	Structural characterization of organized systems of polysaccharides and phospholipids by light scattering spectroscopy and electron microscopy
K. Ogawa (Osaka, Japan), T. Yui (Miyazaki, Japan), K. Nakata (Osaka, Japan), M. Kakuta and A. Misaki (Kobe, Japan)	41	X-ray study of beijeran sodium salts, a new galacturonic acid-containing exo-polysaccharide
E.F. Hounsell (London, UK)	47	Glycosylation targets for drug design
B. Hardy (Oxford, UK)	49	Current topics in carbohydrate modelling: diverse questions, diverse methods
E.R. Caffarena and J.R. Grigera (La Plata, Argentina)	51	Glass transition in aqueous solutions of glucose. Molecular dynamics simulation
S. Abbate, L. Biancardi, G. Longhi (Brescia, Italy) and G. Conti (Novara, Italy)	59	Study of the phase space structure for a two-degrees of freedom model of a symmetric glycosidic linkage
D.A. Brant (Irvine, CA, USA)	67	Polysaccharide and oligosaccharide flexibility
M. Hricovíni, M. Guerrini, G. Torri and B. Casu (Milan, Italy)	69	Motional properties of *E. Coli* polysaccharide K5 in aqueous solution analyzed by NMR relaxation measurements
G. Paradossi, F. Cavalieri and V. Crescenzi (Rome, Italy)	77	^{1}H NMR relaxation study of a chitosan–cyclodextrin network
B. Manunza, S. Deiana, M. Pintore (Sassari, Italy) and C. Gessa (Bologna, Italy)	85	Molecular dynamics study of polygalacturonic acid chains in aqueous solution

B. Manunza, S. Deiana, M. Pintore, G. Delogu (Sassari, Italy) and C. Gessa (Bologna, Italy) 89 A molecular modeling study on the interaction between β-cyclodextrin and synthetic pyretroids

I Announcement

iii Graphical Abstracts

Regular Papers

A. Rydlund (Stockholm, Västerås, Sweden) and O. Dahlman (Stockholm, Sweden) 95 Oligosaccharides obtained by enzymatic hydrolysis of birch kraft pulp xylan: Analysis by capillary zone electrophoresis and mass spectrometry

I. Redmann, M. Pina, B. Guyot (Montpellier, France), P. Blaise, M. Farines (Perpignan, France) and J. Graille (Montpellier, France) 103 Chemoenzymatic synthesis of glucose fatty esters

J. Cai, B.E. Davison, C.R. Ganellin (London, UK), S. Thaisrivongs (Kalamazoo, MI, USA) and K.S. Wibley (London, UK) 109 Potential HIV protease inhibitors: Preparation of di-*N*-alkylated 2-, 6-, and 2,6-amino-deoxy-derivatives of D-glucose by direct displacement and by a novel reductive-alkylation procedure

J. Gervay, T.N. Nguyen and M.J. Hadd (Tucson, AZ, USA) 119 Mechanistic studies on the stereoselective formation of glycosyl iodides: first characterization of β-D-glycosyl iodides

P.B. van Seeventer (Utrecht, The Netherlands), J. Kerékgyártó (Utrecht, The Netherlands; Debrecen, Hungary), J.A.L.M. van Dorst, K.M. Halkes, J.P. Kamerling and J.F.G. Vliegenthart (Utrecht, The Netherlands) 127 Chemical and chemo-enzymatic synthesis of the α-Neu*p*5Ac-(2 → 6)-β-D-Gal*p*NAc-(1 → 4)-β-D-Glc*p*NAc-(1 → 2)-α-D-Man*p* element that is part of N-linked carbohydrate chains of human lutropin

V. Bouchez, I. Stasik, D. Beaupère and R. Uzan (Amiens, France) 139 Regioselective halogenation of pentono-1,4-lactones. Efficient synthesis of 5-chloro- and 5-bromo-5-deoxy derivatives

U. Zähringer (Borstel, Germany), H. Rettenmaier (Ludwigshafen, Germany), H. Moll (Borstel, Germany), S.N. Senchenkova (Moscow, Russian Federation) and Yu.A. Knirel (Borstel, Germany; Moscow, Russian Federation) 143 Structure of a new 6-deoxy-α-D-talan from *Burkholderia* (*Pseudomonas*) *plantarii* strain DSM 6535, which is different from the O-chain of the lipopolysaccharide

B. Evers (Hamburg, Germany), M. Petříček (Prague, Czech Republic) and J. Thiem (Hamburg, Germany) 153 Specificity of amylases and cyclodextrin-glucanotransferase in reactions with 2-deoxy-maltooligosaccharides

A. Lubineau, K. Basset-Carpentier and C. Augé (Orsay, France) 161 Porcine liver (2 → 3)-α-sialyltransferase: substrate specificity studies and application of the immobilized enzyme to the synthesis of various sialylated oligosaccharide sequences

Notes

C.P.J. Glaudemans, C.E. Miller and E.M. Nashed (Bethesda, MD, USA) 169 The measurement of affinity between ligand and antibody using ligand-induced antibody fluorescence change

W.-Y. Wu, B. Jin, D.C.M. Kong and M. von Itzstein (Parkville, Australia) 171 A facile synthesis of a useful 5-*N*-substituted-3,5-dideoxy-D-*glycero*-D-*galacto*-2-non-ulosonic acid from 2-acetamido-2-deoxy-D-glucose

A. Morère, C. Menut, C. Vidil, P. Skaanderup, J. Thorsen, J.-P. Roque and J.-L. Montero (Montpellier, France) 175 The benzyloxycarbonyl group: An alternative protective group in the mannose series

A. Oosterveld (Wageningen, The Netherlands), J.H. Grabber (Madison, WI, USA), G. Beldman (Wageningen, The Netherlands), J. Ralph (Madison, WI, USA) and A.G.J. Voragen (Wageningen, The Netherlands) 179 Formation of ferulic acid dehydrodimers through oxidative cross-linking of sugar beet pectin

Z.G. Tóth, I.F. Pelyvás, C. Szegedi, P. Benke, E. Magyar, T. Miklovicz, G. Batta and F. Sztaricskai (Debrecen, Hungary) 183 Synthesis of new anhydro and branched-chain cyclitols

C. Karlsson, P.-E. Jansson (Huddinge, Sweden) and R. Wollin (Stockholm, Sweden) 191 Structure of the O-polysaccharide from the LPS of a *Hafnia alvei* strain isolated from a patient with suspect yersinosis

iii Graphical Abstracts

Regular Papers

W.S. York, S. Hantus, P. Albersheim and A.G. Darvill (Athens, GA, USA) 199 Determination of the absolute configuration of monosaccharides by ^{1}H NMR spectroscopy of their per-*O*-(*S*)-2-methylbutyrate derivatives

H. Ando, H. Ishida, M. Kiso and A. Hasegawa (Gifu, Japan) 207 A synthetic approach to the c-series gangliosides containing sialyl-α(2 → 8)sialyl-α(2 → 8)sialic acid: Synthesis of ganglioside GT4, α(2 → 6)GT4 and GT3

J.-l. Jane, K.-s. Wong and A.E. McPherson (Ames, IA, USA) 219 Branch-structure difference in starches of A- and B-type X-ray patterns revealed by their Naegeli dextrins

R. Panikkar and D.J. Brasch (Dunedin, New Zealand) 229 Biosynthetic implications of NMR analyses of alginate homo- and heteropolymers from New Zealand brown seaweeds

K.-S. Shin, H. Kiyohara, T. Matsumoto and H. Yamada (Tokyo, Japan) 239 Rhamnogalacturonan II from the leaves of *Panax ginseng* C.A. Meyer as a macrophage Fc receptor expression-enhancing polysaccharide

A.E. Frankel, T. Fu, C. Burbage, E. Tagge, B. Harris, J. Vesely and M.C. Willingham (Charleston, SC, USA) 251 Lectin-deficient ricin toxin intoxicates cells bearing the D-mannose receptor

Y. Kameda, K. Kawashima, M. Takeuchi, K. Ikeda, N. Asano and K. Matsui (Kanazawa city, Japan) 259 Preparation and biological activity of manno- and galacto-validamines, new 5a-carba-glycosylamines as α-glycosidase inhibitors

Notes

U. Büttner, F. Gerum and T. Severin (München, Germany) 265 Formation of α-amino-acid amides and α-hydroxy-acid amides by degradation of sugars with primary amines

V. Moreau, J.Chr. Norrild and H. Driguez (Grenoble, France) 271 Synthesis of methyl 4-thio-β-cellobioside. A reinvestigation

S.J. Angyal (Sydney, Australia) 279 A short note on the epimerization of aldoses

Y. Nishiyama, N. Nishimura, N. Kuroyanagi and I. Maeba (Nagoya, Japan) 283 Synthesis of 3-β-D-ribofuranosyl-1*H*-pyrazole-4-carboxamide

iii Graphical Abstracts

Regular Papers

D. Bailey, M.J. Davies, F.H. Routier, C. Bauer, J. Feeney and E.F. Hounsell (London, UK) 289 ^{1}H NMR analysis of novel sialylated and fucosylated lactose-based oligosaccharides having linear GlcNAc(β1–6)Gal and Neu5Ac(α2–6)GlcNAc sequences

J.S. Kozlowski, C.H. Marzabadi, N.P. Rath and C.D. Spilling (St. Louis, MO, USA) 301 Chemistry of glucal halohydrins (II): An unusual protecting group effect in the competitive formation of formyl furanosides and methyl glycosides

P.A.S. Mourão, M.S.G. Pavão (Rio de Janeiro, Brazil), B. Mulloy (Potters Bar, UK) and R. Wait (Salisbury, UK) 315 Chondroitin ABC lyase digestion of an ascidian dermatan sulfate. Occurrence of unusual 6-*O*-sulfo-2-acetamido-2-deoxy-3-*O*-(2-*O*-sulfo-α-L-idopyranosyluronic acid)-β-D-galactose units

S.F. Osman, W.F. Fett, P. Irwin (Wyndmoor, PA, USA), P. Cescutti (Trieste, Italy), J.N. Brouillette and J.V. O'Connor (Wyndmoor, PA, USA) 323 The structure of the exopolysaccharide of *Pseudomonas fluorescens* strain H13

J. Zhang and P. Kováč (Bethesda, MD, USA) 329 Synthesis of methyl α-glycosides of some higher oligosaccharide fragments of the O-antigen of *Vibrio cholerae* O1, serotype Inaba and Ogawa

J.-N. Bertho (Rennes, France), A. Coué, D.F. Ewing, J.W. Goodby, P. Letellier, G. Mackenzie (Hull, UK) and D. Plusquellec (Rennes, France) 341 Novel sugar bola-amphiphiles with a pseudo macrocyclic structure

Notes

W. Liao and D. Lu (Beijing, PR China)	347	Convergent synthesis of an octasaccharide acceptor corresponding to the reducing terminus of mycobacterial 3-*O*-methylmannose polysaccharide (MMP)
P.A. Kroon, M.T. Garcia Conesa, I.J. Colquhoun and G. Williamson (Norwich, UK)	351	Process for the isolation of preparative quantities of [2-*O*-(*trans*-feruloyl)-α-L-arabinofuranosyl]-(1 → 5)-L-arabinofuranose from sugarbeet
J. Ning and F. Kong (Beijing, PR China)	355	Synthesis and glycosidic coupling reaction of substituted 2,6-dioxabicyclo[3.1.0]hexanes: 1,2-anhydro-3,5-di-*O*-benzyl-α-D-ribofuranose
J. Jindrich (Praha, Czech Republic), K. Harata (Ibaraki, Japan), B. Lindberg, J. Pitha and P. Seffers (Stockholm, Sweden)	361	Synthesis of $2^1,3^1$-*O*-(propane-1,2-diyl)- and $2^1,3^1$-*O*-(3-hydroxypropane-1,2-diyl)-cyclomaltoheptaose
J.O. Hoberg (Golden, CO, USA)	365	Synthesis and chemistry of 4,6-*O*-di-(*tert*-butyl)silanediyl-D-glucal
P.B. van Seeventer, J.A.L.M. van Dorst, J.F. Siemerink, J.P. Kamerling and J.F.G. Vliegenthart (Utrecht, The Netherlands)	369	Thiol addition to protected allyl glycosides: an improved method for the preparation of spacer-arm glycosides
R.R. Diaz, C.R. Melgarejo, I.I. Cubero and M.T. Plaza López-Espinosa (Granada, Spain)	375	Synthesis of furanose glycals from furanose 1,2-diols and their cyclic thiocarbonate esters
	III	Author index
	V	Subject index
	XI	Contents